

Reply to the Comment [1] on “Scaling Laws for a System with Long-Range Interactions within Tsallis Statistics” [2]

R. Salazar and R. Toral

Departament de Física, Universitat de les Illes Balears

Instituto Mediterráneo de Estudios Avanzados (IMEDEA, UIB-CSIC)

07071 Palma de Mallorca, Spain.

The fact that mean field theory is appropriate to describe an Ising model with long-range interactions has been already shown by Cannas and Tamarit [3]. Although not explicated in our Letter, we have used periodic boundary conditions in all our simulations, such that the maximum possible distance between two lattice sites is $L/2$.

We make no specific comments about the validity (or lack of validity) of Boltzmann-Gibbs statistical mechanics to describe long-range systems. Our paper does nothing but to compare the scaling functions derived from Boltzmann-Gibbs statistics from those derived from the use of Tsallis entropy.

-
- [1] B.P. Vollmayr-Lee and E. Luijten, to appear in *Phys. Rev. Lett.*, preprint cond-mat/0005444.
 - [2] R. Salazar and R. Toral, *Phys. Rev. Lett.* **83** (1999) 4233.
 - [3] S.A. Cannas and F.A. Tamarit, *Phys. Rev.* **B54**, R12661 (1996). S.A. Cannas, A. de Magalhaes and F.A. Tamarit, preprint cond-mat/9906340.